

AMENDMENTS TO THE SPECIFICATION

The Specification as amended below show added text with underlining and deleted text with ~~strikethrough~~.

Please REPLACE paragraph [0023] with the following amended paragraph:

[0023] The 2^n level quantizing method will be described in more detail below. If the samples $r(k)$ in the above-described Equation (1) are substituted with 2^n level quantized samples $q(k)$, the following equation is obtained:

$$\Lambda(n) = \sum_{k=1}^{n+L} q(k)q^*(k-N) \quad \dots (2)_{\pm}$$

Please REPLACE paragraph [0024] with the following amended paragraph:

[0024] If a sample $\max r(k)$ having a largest value among the samples $r(k)$ is 2^n , and the other samples $r(k)$ are proportionally magnified or scaled, the 2^n level quantized samples $q(k)$ can be represented by a quantization function Q_L as follows:

$$q(k) = Q_L \left[\frac{2^n r(k)}{\max r(k)} \right] \quad \dots (3)_{\pm}$$

Please REPLACE paragraph [0025] with the following amended paragraph:

[0025] The term $Q_L[x]$ in Equation (3) represents a complex quantization to quantize the scaled sample values into levels of 2^i in accordance with the following equation:

$$Q_L[x] \cong Q[\operatorname{Re}\{x\}] + jQ[\operatorname{Im}\{x\}] \quad \dots (4)_{\pm}$$

Please REPLACE paragraph [0026] with the following amended paragraph:

[0026] The term $Q_L[x]$ can also be represented as follows:

$$Q_L(x) \cong \begin{cases} 2^{\lceil \log_2 x \rceil}, & x > 0 \\ 0, & x = 0 \end{cases} \quad \dots (5)_{\pm}$$

Please REPLACE paragraph [0028] with the following amended paragraph:

[0028] If the shifter is used instead of the multiplier, Equation (1) can be represented as follows:

$$\Lambda(n) = \sum_{k=1}^{n+L} \{q(k) \ll l(k-N)\} \quad \dots (6)$$

$$l(k-N) = \log_2 q^*(k-N) \quad \dots (7)$$

Please DELETE the paragraph number [0031] between paragraphs [0030] and [0032].